



• **FACT SHEET No. 7**

Pain After Surgery in Children and Infants

Children have at least the same amount of pain from surgery as adults do, although it may not last as long because children usually heal more quickly. Children deserve protection, cannot advocate for themselves, and often may not complain of pain, so they need special consideration and pain assessment. All health professionals should know how to recognize, assess, and treat pain in children. [Finley 2005]

Untreated acute pain can result in chronic pain in children and adolescents [Batoz 2016], and failure to prevent pain in newborns can cause lifelong adverse effects, such as increased pain sensitivity. [Vinnall 2014]

Surgical pain should be prevented whenever possible. Avoid unnecessary procedures and plan the management before the surgery. Assess pain using age-appropriate validated tools. Develop standard protocols that can be adapted to individual patients, so that when surgery is required, a combination of medications are available to provide the best possible analgesia with the lowest risk and side-effects, along with non-pharmacological pain control techniques. [AAP 2001] [APAGBI 2012]

Local/Regional Techniques

Local anesthetics can block pain nerves before, during, and after the surgical procedure.

- Local anesthetic infiltration at surgical site
- Local nerve block
- Plexus or nerve sheath block
- Neuraxial block (epidural, spinal)

Medications

Most medications are safe in children and infants, if the dosing is adjusted for weight and for the metabolic differences in newborns and infants. All of the following drugs may be used with appropriate



© Copyright 2017 International Association for the Study of Pain. All rights reserved.

IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.

monitoring to treat pain and to reduce nerve sensitization, depending on the patient and type of surgery or injury.

- Paracetamol/Acetaminophen
- Non-steroidal anti-inflammatory drugs
- Opioids
- Gabapentin
- Ketamine

Psychological and Physical Approaches

Psychological and physical techniques are very important, and are also inexpensive and safe. They should be available for all patients.

- Preparation and explanation
- Distraction, imaging, and relaxation (including deep breathing) [Davidson 2016]
- Positioning and early gentle movement

Parents' Role

- Advocating for their child's care
- Providing distraction, support, and comfort

Professionals' Role

- Be aware of the child's pain.
- Think of pain prevention before it happens. Do not do unnecessary procedures
- Recognize, assess, prevent, treat

REFERENCES

Finley GA, Franck LS, Grunau RE, von Baeyer CL. Why children's pain matters. *Pain: Clinical Updates* 2005, Sep;13(4):1-6.

Batoz H, Semjen F, Bordes-Demolis M, Bénard A, Nouette-Gaulain K. Chronic postsurgical pain in children: prevalence and risk factors. A prospective observational study. *Br J Anaesth* 2016; 117(4): 489-96

Vinall J, Grunau RE. Impact of repeated procedural pain-related stress in infants born very preterm. *Pediatr Res* 2014, May;75(5):584-7.

American Academy of Pediatrics. The assessment and management of acute pain in infants, children, and adolescents. *Pediatrics* 2001, Sep 1;108(3):793-7.

Association of Paediatric Anaesthetists of Great Britain and Ireland. Good practice in postoperative and procedural pain management, 2nd edition. *Paediatr Anaesth* 2012, Jul;22 Suppl 1:1-79.

Davidson F, Snow S, Hayden JA, Chorney J. Psychological interventions in managing postoperative pain in children: A systematic review. *PAIN* 2016; 157(9): 1872-86



© Copyright 2017 International Association for the Study of Pain. All rights reserved.

IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.

AUTHOR

G. Allen Finley, MD FRCPC FAAP
Professor of Anesthesia and Psychology, Dalhousie University
Dr. Stewart Wenning Chair in Pediatric Pain Management
Director, Centre for Pediatric Pain Research, IWK Health Centre
Halifax, Canada

REVIEWERS

Supranee Niruthisard, MD
Associate Professor in Anesthesiology
Department of Anesthesiology, Chulalongkorn University
King Chulalongkorn Memorial Hospital
Bangkok, Thailand

Jill Chorney, PhD, RPsych
Associate Professor of Anesthesia & Psychology
Dalhousie University
Halifax, Canada

A. Stuart Wright, MD PhD FRCPC
Assistant Professor of Anesthesia
Dalhousie University
Halifax, Canada

About the International Association for the Study of Pain®

IASP is the leading professional forum for science, practice, and education in the field of pain. [Membership is open to all professionals](#) involved in research, diagnosis, or treatment of pain. IASP has more than 7,000 members in 133 countries, 90 national chapters, and 20 Special Interest Groups.

As part of the Global Year Against Pain After Surgery, IASP offers a series of Fact Sheets that cover specific topics related to postsurgical pain. These documents have been translated into multiple languages and are available for free download. Visit www.iasp-pain.org/globalyear for more information.



© Copyright 2017 International Association for the Study of Pain. All rights reserved.

IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.